RECEIVED CENTRAL FAX CENTER DEC 1 9 2006

## IN THE CLAIMS:

- (Original) A polymerization process comprising contacting one or more monomer(s), one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor.
- (Original) The process of claim 1, wherein one or more hydrofluorocarbon(s) is represented by the formula: C<sub>x</sub>H<sub>y</sub>F<sub>z</sub> wherein x is an integer from 1 to 40 and y and z are integers of one or more.
- 3. (Original) The process of claim 2, wherein x is from 1 to 10.
- 4. (Original) The process of claim 2, wherein x is from 1 to 6.
- 5. (Original) The process of claim 2, wherein x is from 1 to 3.
- (Original) The process of claim 1, wherein the one or more hydrofluorocarbon(s) is 6. independently selected from the group consisting of fluoromethane; difluoromethane; trifluoromethane: fluoroethane; 1,1-difluoroethane; 1,2-difluoroethane; 1,1,1-1,1,2,2trifluoroethane; 1,1,2-trifluoroethane; 1,1,1,2-tetrafluoroethane; tetrafluoroethane; 1,1,1,2,2-pentafluoroethane; 1-fluoropropane; 2-fluoropropane; 1,1difluoropropane; 1,2-difluoropropane; 1,3-difluoropropane; 2,2-difluoropropane; 1,1,1-trifluoropropane; 1,2,2-1,1,2-trifluoropropane; 1,1,3-trifluoropropane; 1,1,1,2-tetrafluoropropane; 1,1,1,3trifluoropropane; 1.2.3-trifluoropropane; tetrafluoropropane; 1,1,2,2-tetrafluoropropane; 1,1,2,3-tetrafluoropropane; 1,1,3,3-1,1,1,2,2-pentafluoropropane; tetrafluoropropane; 1,2,2,3-tetrafluoropropane; 1,1,1,2,3-pentafluoropropane; 1,1,1,3,3-pentafluoropropane; 1,1,2,2,3-1,1,2,3,3-pentafluoropropane; 1,1,1,2,2,3-hexafluoropropane; pentafluoropropane; 1,1,1,3,3,3-hexafluoropropane; 1.1.1.2.2.3.3-1,1,1,2,3,3-hexafluoropropane; 1,1,1,2,3,3,3-heptafluoropropane; 2heptafluoropropane; 1-fluorobutane; fluorobutane; 1,1-difluorobutane; 1,2-difluorobutane; 1,3-difluorobutane; 1,4-

> difluorobutane; 2,2-difluorobutane; 2,3-difluorobutane; 1,1,1-trifluorobutane; 1,1,2trifluorobutane; 1,1,3-trifluorobutane; 1,1,4-trifluorobutane; 1,2,2-trifluorobutane; 1,1,1,2-1,3,3-trifluorobutane; 2,2,3-trifluorobutane; 1.2.3-trifluorobutane: 1,1,2,2-1,1,1,4-tetrafluorobutanc; tetrafluorobutane; 1,1,1,3-tetrafluorobutane; 1,1,2,3-tetrafluorobutane; 1,1,2,4-tetrafluorobutane; 1,1,3,3tetrafluorobutane; 1,1,3,4-tetrafluorobutane; 1,1,4,4-tetrafluorobutane; 1,2,2,3tetrafluorobutane: 1,2,2,4-tetrafluorobutane; 1,2,3,3-tetrafluorobutane; 1,2,3,4tetrafluorobutane; tetrafluorobutane; 2,2,3,3-tetrafluorobutane; 1,1,1,2,2-pentafluorobutane; 1,1,1,2,3-1,1,1,2,4-pentafluorobutane; 1,1,1,3,3-pentafluorobutane; pentafluorobutane; 1,1,2,2,3-1,1,1,4,4-pentafluorobutane; 1,1,3,4-pentafluorobutane; 1,1,2,3,3-pentafluorobutane; 1,1,2,2,4-pentafluorobutane; pentafluorobutane; 1,1,3,3,4-pentafluorobutane; 1,2,2,3,3-1,1,2,4,4-pentafluorobutane; 1,1,1,2,2,3-hexafluorobutane; pentafluorobutane; 1,2,2,3,4-pentafluorobutane; 1,1,1,2,3,3-hexafluorobutanc, 1.1.1.2.3.4-1,1,1,2,2,4-hexafluorobutane; 1,1,1,2,4,4-hexafluorobutane; 1,1,1,3,3,4-hexafluorobutane; hexafluorobutane; 1,1,2,2,3,3-1,1,4,4,4-hexafluorobutane; 1,1,1,3,4,4-hexafluorobutane; 1.1.2.2.4.4-hexafluorobutane; 1,1,2,2,3,4-hexafluorobutane; hexafluorobutane; 1,2,2,3,3,4-1,1,2,3,3,4-hexafluorobutane; 1,1,2.3,4,4-hexafluorobutane; hexafluorobutane; 1,1,1,2,2,3,3-heptafluorobutane; 1,1,1,2,2,4,4-heptafluorobutane; 1,1,1,2,2,3,4-heptafluorobutane; 1,1,1,2,3,3,4-heptafluorobutane; 1,1,1,2,3,4,4heptafluorobutane; 1,1,1,2,4,4,4-heptafluorobutane; 1,1,1,3,3,4,4-heptafluorobutane; 1,1,1,2,2,3,3,4-octafluorobutane; 1,1,1,2,2,3,4,4-octafluorobutane; octafluorobutane; 1,1,1,2,2,4,4,4-octafluorobutane; 1,1,1,2,3,4,4,4-octafluorobutane; 1.1.1.2.2.3.3.4.4-nonafluorobutane; 1.1.1.2.2.3.4.4.4-nonafluorobutane; 1-fluoro-2methylpropane; 1,1-difluoro-2-methylpropane; 1,3-difluoro-2-methylpropane; 1,1,1-1,1,3-trifluoro-2-methylpropane; 1,3-difluoro-2trifluoro-2-methylpropane; (fluoromethyl)propanc; 1,1,1,3-tetrafluoro-2-methylpropane; 1,1,3,3-tetrafluoro-2-1,1,3-trifluoro-2-(fluoromethyl)propane; 1,1,1,3,3-pentafluoro-2methylpropane: methylpropane; 1.1,3,3-tetrafluoro-2-(fluoromethyl)propane; 1,1,1,3-tetrafluoro-2-1.1-difluorocyclobutane; 1,2fluorocyclobutane; (fluoromethyl)propane; difluorocyclobutane; 1.3-difluorocyclobutane; 1,1,2-trifluorocyclobutane; 1,1,3-1,1,2,2-tetrafluorocyclobutane; trifluorocyclobutane; 1,2,3-trifluorocyclobutane;

> 1,1,2,2,3-pentafluorocyclobutane; 1,1,2,3,3-1,1,3,3-tetrafluorocyclobutane; 1,1,2,2,3,3-hexafluorocyclobutane; pentafluorocyclobutane; 1, 1, 2, 2, 3, 4hexafluorocyclobutane; 1,1,2,3,3,4-hexafluorocyclobutane; 1,1,2,2,3,3,4heptafluorocyclobutane; vinyl fluoride; 1,1-difluoroethene; 1,2-difluoroethene; 1,1,2trifluoroethene; 1-fluoropropene, 1,1-difluoropropene; 1,2-difluoropropene; 1,3difluoropropene; 2,3-difluoropropene; 3,3-difluoropropene; 1,1,2-trifluoropropene; 1,1,3-trifluoropropene; 1,2,3-trifluoropropene; 1,3,3-trifluoropropene; trifluoropropene; 3,3,3-trifluoropropene; 1-fluoro-1-butene; 2-fluoro-1-butene; 3fluoro-1-butene; 4-fluoro-1-butene; 1,1-difluoro-1-butene; 1,2-difluoro-1-butene; 1,3difluoropropene; 1,4-difluoro-1-butene; 2,3-difluoro-1-butene; 2,4-difluoro-1-butene; 3,3-difluoro-1-butene; 3,4-difluoro-1-butene; 4,4-difluoro-1-butene; 1,1,2-trifluoro-1butene; 1,1,3-trifluoro-1-butene; 1,1,4-trifluoro-1-butene; 1,2,3-trifluoro-1-butene; 1,2,4-trifluoro-1-butene; 1,3,3-trifluoro-1-butene; 1,3,4-trifluoro-1-butene; 1,4,4trifluoro-1-butene; 2,3,3-trifluoro-1-butene; 2,3,4-trifluoro-1-butene; 2,4,4-trifluoro-1butene; 3,3,4-trifluoro-1-butene; 3,4,4-trifluoro-1-butene; 4,4,4-trifluoro-1-butene; 1,1,2,4-tetrafluoro-1-butene; 1,1,3,3-tetrafluoro-1-1,1,2,3-tetrafluoro-1-butene; butene: 1.1.3.4-tetrafluoro-1-butene: 1.1.4.4-tetrafluoro-1-butene; 1,2,3,3-tetrafluoro-1.2.4.4-tetrafluoro-1-butene; 1.3.3.4-1-butene: 1,2,3,4-tetrafluoro-1-butene; 1.4.4.4-tetrafluoro-1-butene; tetrafluoro-1-butene; 1,3,4,4-tetrafluoro-1-butene; 2,4,4,4-tetrafluoro-1-2.3.3.4-tetrafluoro-1-butene: 2,3,4,4-tetrafluoro-1-butene; 3,3,4,4-tetrafluoro-1-butene; 3,4,4,4-tetrafluoro-1-butene; 1,1,2,3,3butene; pentafluoro-1-butene; 1,1,2,3,4-pentafluoro-1-butene; 1,1,2,4,4-pentafluoro-1-butene; 1,1,3,3,4-pentafluoro-1-butene; 1,1,3,4,4-pentafluoro-1-butene; 1,1,4,4,4-pentafluoro-1-butene; 1,2,3,3,4-pentafluoro-1-butene; 1,2,3,4,4-pentafluoro-1-butene; 1,2,4,4,4pentafluoro-1-butene: 2,3,3,4,4-pentafluoro-1-butene; 2,3,4,4,4-pentafluoro-1-butene; 3,3,4,4,4-pentafluoro-1-butene; 1,1,2,3,3,4-hexafluoro-1-butene; 1,1,2,3,4,4-1,1,2,4,4,4-hexafluoro-1-butene; 1,2,3,3,4,4-hexafluoro-1hexafluoro-1-butene; 1,2,3,4,4,4-hexafluoro-1-butene; 2,3,3,4,4,4-hexafluoro-1-butene; butene; 1,1,2,3,3,4,4-heptafluoro-1-butene; 1,1,2,3,4,4,4-heptafluoro-1-butene; 1,1,3,3,4,4,4heptafluoro-1-butene; 1,2,3,3,4,4,4-heptafluoro-1-butene; 1-fluoro-2-butene; 2-fluoro-2-butene; 1,1-difluoro-2-butene; 1,2-difluoro-2-butene; 1,4difluoro-2-butene; 2,3-difluro-2-butene; 1,1,1-trifluoro-2-butene; 1,1,2-trifluoro-2-

> butene; 1,1,3-trifluoro-2-butene; 1,1,4-trifluoro-2-butene; 1,2,3-trifluoro-2-butene; 1,2,4-trifluoro-2-butene; 1,1,1,2-tetrafluoro-2-butene; 1,1,1,3-tetrafluoro-2-butene; 1,1,2,3-tetrafluoro-2-butene; 1,1,2,4-tetrafluoro-2-1,1,1,4-tetrafluoro-2-butene; 1,2,3,4-tetrafluoro-2-butene; 1,1,1,2,3-pentafluoro-2-butene; 1,1,1,2,4pentafluoro-2-butene; 1,1,1,3,4-pentafluoro-2-butene; 1,1,1,4,4-pentafluoro-2-butene; 1,1,2,4,4-pentafluoro-2-butene; 1,1,1,2,3,4-1,1,2,3,4-pentafluoro-2-butene; 1,1,1,3,4,4-hexafluoro-2-1,1,1,2,4,4-hexafluoro-2-butene; hexafluoro-2-butene; 1,1,1,4,4,4-hexafluoro-2-butene; 1,1,2,3,4,4-hexafluoro-2-butene; butene; 1,1,1,2,3,4,4-heptafluoro-2-butene; 1,1,1,2,4,4,4-heptafluoro-2-butene; and mixtures thereof.

- 7. (Original) The process of claim 1, wherein the one or more hydrofluorocarbon(s) is independently selected from the group consisting of fluoromethane, difluoromethane, trifluoromethane, 1,1-difluoroethane, 1,1-trifluoroethane, 1,1,2-tetrafluoroethane, and mixtures thereof.
- (Previously presented) The process of claim 1, wherein the diluent comprises from 15
   to 100 volume % HFC based upon the total volume of the diluent.
- (Original) The process of claim 8, wherein the diluent comprises from 20 to 100 volume % HFC based upon the total volume of the diluent.
- (Original) The process of claim 9, wherein the diluent comprises from 25 to 100 volume % HFC based upon the total volume of the diluent.
- 11. (Previously presented) The process of claim 1, wherein the diluent further comprises a hydrocarbon, a non-reactive olefin, and/or an inert gas.
- 12. (Currently amended) The process of <u>claim 1</u>, wherein the <u>diluent further comprises</u> elaim 11, wherein the <u>hydrocarbon</u> is a halogenated hydrocarbon other than an HFC.

- 13. (Currently amended) The process of claim 12, wherein the halogenated hydrocarbon is claim 1, wherein the diluent further comprises methyl chloride.
- 14. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>4</sub>; wherein M is a Group 4, 5, or 14 metal; and each X is a halogen.
- 15. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MR<sub>n</sub>X<sub>4-n</sub>; wherein M is Group 4, 5, or 14 metal; each R is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4; and each X is a halogen.
- (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula M(RO)<sub>n</sub>R'<sub>m</sub>X<sub>4-(m+n)</sub>. wherein M is Group 4, 5, or 14 metal; each RO is a monovalent C<sub>1</sub> to C<sub>30</sub> hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, arylalkoxy, alkylaryloxy radicals; each R' is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4; m is an integer from 0 to 4, wherein the sum of n and m is not more than 4; and each X is a halogen.
- 17. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula M(RC=OO)<sub>n</sub>R'<sub>m</sub>X<sub>4-(m+n)</sub>; wherein M is Group 4, 5, or 14 metal;

each RC=OO is a monovalent  $C_2$  to  $C_{30}$  hydrocarbacyl radical independently selected from the group consisting of an alkacyloxy, arylacyloxy, arylacyloxy, alkylarylacyloxy radicals; each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals;

n is an integer from 0 to 4;
m is an integer from 0 to 4, wherein the sum of n and m is not more than 4; and

m is an integer from 0 to 4, wherein the sum of n and m is not more than 4; and each X is a halogen.

- 18. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MOX<sub>3</sub>; wherein M is a Group 5 metal; and each X is a halogen.
- 19. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>3</sub>; wherein M is a Group 13 metal; and each X is a halogen.
- 20. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MR<sub>n</sub>X<sub>3-n</sub>; wherein M is a Group 13 metal; each R is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 1 to 3; and each X is a halogen.
- 21. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula M(RO)<sub>n</sub>R'<sub>m</sub>X<sub>3-(m+n)</sub>: wherein M is a Group 13 metal; each RO is a monovalent C<sub>1</sub> to C<sub>30</sub> hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, arylakoxy, alkylaryloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 3; m is an integer from 0 to 3, wherein the sum of n and m is from 1 to 3; and each X is a halogen.

22. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula M(RC=OO)<sub>n</sub>R'<sub>m</sub>X<sub>3-(m+n)</sub>; wherein M is a Group 13 metal; each RC=OO is a monovalent hydrocarbacyl radical independently selected from the

each RC=OO is a monovalent hydrocarbacyl radical independently selected from the group independently selected from the  $C_2$  to  $C_{30}$  group consisting of an alkacyloxy, arylacyloxy, arylacyloxy, alkylarylacyloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 3;

m is a integer from 0 to 3, wherein the sum of n and m is from 1 to 3; and each X is a halogen.

- 23. (Currently amended) A polymerization process comprising contacting one or more isoplefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor, The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>y</sub>, wherein M is a Group 15 metal; each X is a halogen; and y is 3, 4 or 5.
- 24. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of elaim-1, wherein the one or more Lewis acid(s) is represented by the formula:

  MR<sub>n</sub>X<sub>y-n:</sub>

  wherein M is a Group 15 metal;

each R is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4;

y is 3, 4 or 5, wherein n is less than y; wherein the sum of n and m is less than y each X is a halogen.

25. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of claim 1, wherein the one or more Lewis acid(s) is represented by the formula M(RO)<sub>0</sub>R'<sub>m</sub>X<sub>v-(m+n)</sub>:

wherein M is a Group 15 metal,

each RO is a monovalent  $C_1$  to  $C_{30}$  hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, aryloxy, alkylaryloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals;

n is an integer from 0 to 4;

m is an integer from 0 to 4;

y is 3, 4 or 5, wherein the sum of n and m is less than y; and each X is a halogen.

26. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of elaim—I, wherein the one or more Lewis acid(s) is represented by the formula M(RC=OO)<sub>n</sub>R'<sub>m</sub>X<sub>y-(m+n)</sub>;

wherein M is a Group 15 metal;

each RC=OO is a monovalent  $C_2$  to  $C_{30}$  hydrocarbacyloxy radical independently selected from the group consisting of an alkacyloxy, arylacyloxy, arylacyloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals;

> n is an integer from 0 to 4; m is an integer from 0 to 4; y is 3, 4 or 5, wherein the sum of n and m is less than y; and each X is a halogen.

(Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) 27. is independently selected from the group consisting of titanium tetrachloride, titanium tetrabromide, vanadium tetrachloride, tin tetrachloride, zirconium tetrachloride, titanium bromide trichloride, titanium dibromide dichloride, vanadium bromide trichloride, tin chloride trifluoride, benzyltitanium trichloride, dibenzyltitanium dibromide, dibenzylzirconium dichloride. benzylzirconium trichloride, methyltitanium trichloride, dimethyltitanium difluoride, dimethyltin dichloride, n-butoxytitanium methoxytitanium trichloride. trichloride, phenylvanadium phenoxytitanium tribromide, trichloride, di(isopropoxy)titanium dichloride, phenylmethoxyzirconium trifluoride, methyl methoxytitanium dichloride, methyl methoxytin dichloride, benzyl isopropoxyvanadium dichloride, acetoxytitanium benzoyloxytitanium trifluoride, tribromide, trichloride, benzoylzirconium dichloride, trichloride, methyl acetoxytitanium benzyl isopropoyloxytin benzoyloxyvanadium chloride, vanadium oxytrichloride, aluminum trichloride, boron trifluoride, gallium trichloride, indium trifluoride, ethylaluminum dichloride, methylaluminum dichloride, benzylaluminum dichloride, isobutylgallium dichloride, dimethylaluminum chloride. ethylaluminum diethylaluminum chloride. trimethylaluminum, methylaluminum sesquichloride sesquichloride, tricthylaluminum, methoxyaluminum dichloride, ethoxyaluminum dichloride, 2,6-ditert-butylphenoxyaluminum dichloride, methoxy methylaluminum chloride, 2,6-ditert-butylphenoxy methylaluminum chloride, isopropoxygallium dichloride, phenoxy fluoride, acetoxyaluminum dichloride. benzoyloxyaluminum methylindium dibromide, benzoyloxygallium difluoride, methyl acetoxyaluminum chloride, isopropoyloxyindium trichloride, antimony hexachloride, antimony hexafluoride, arsenic pentafluoride, antimony chloride pentafluoride, arsenic trifluoride, bismuth tetraphenylantimony chloride, tetrachloride. trichloride arsenic fluoride tetrachloromethoxyantimony, triphenylantimony dichloride.

dimethoxytrichloroantimony, dichloromethoxyarsine, chlorodimethoxyarsine, difluoromethoxyarsine, acetatotetrachloroantimony, (benzoato) tetrachloroantimony, and bismuth acetate chloride.

- 28. (Previously presented) The process of claim 1, wherein the one or more Lewis acid(s) is independently selected from the group consisting of aluminum trichloride, aluminum tribromide, ethylaluminum dichloride, ethylaluminum sesquichloride, diethylaluminum chloride, methylaluminum dichloride, methylaluminum sesquichloride, dimethylaluminum chloride, boron trifluoride, and titanium tetrachloride.
- 29. (Previously presented) The process of claim 1, wherein the one or more initiator(s) comprise a hydrogen halide, a carboxylic acid, a carboxylic acid halide, a sulfonic acid, an alcohol, a phenol, a polymeric halide, a tertiary alkyl halide, a tertiary aralkyl halide, a tertiary alkyl ester, a tertiary aralkyl ester, a tertiary aralkyl ester, a tertiary aralkyl ether, a tertiary aralkyl ether, an alkyl halide, an aryl halide, an alkylaryl halide or an arylalkylacid halide.
- 30. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor, The process of claim 1, wherein the one or more initiator(s) is independently selected from the group consisting of HCl, H<sub>2</sub>O, methanol, (CH<sub>3</sub>)<sub>3</sub>CCl, C<sub>6</sub>H<sub>5</sub>C(CH<sub>3</sub>)<sub>2</sub>Cl, (2-Chloro-2,4,4-trimethylpentane) and 2-chloro-2-methylpropane.
- 31. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor, The process of elaim-1, wherein the one or more initiator(s) is independently selected from the group consisting of hydrogen chloride, hydrogen bromide, hydrogen iodide, acetic acid, propanole acid, butanole acid; cinnamic acid, benzoic acid, 1-chloroacetic acid, dichloroacetic acid, trichloroacetic acid, trifluoroacetic acid, p-chlorobenzoic acid, p-

fluorobenzoic acid, acetyl chloride, acetyl bromide, cinnamyl chloride, benzoyl chloride, benzoyl bromide, trichloroacetylchloride, trifluoroacetylchloride, p-fluorobenzoylchloride, methanesulfonic acid, trifluoromethanesulfonic acid, trichloromethanesulfonic acid, methanesulfonic acid, methanesulfonyl chloride, methanesulfonyl bromide, trichloromethanesulfonyl chloride, trifluoromethanesulfonyl chloride, p-toluenesulfonyl chloride, methanol, ethanol, propanol, 2-propanol, 2-methylpropan-2-ol, cyclohexanol, benzyl alcohol, phenol, 2-methylphenol, 2,6-dimethylphenol, p-chlorophenol, p-fluorophenol, 2,3,4,5,6-pentafluorophenol, and 2-hydroxynaphthalene.

(Currently amended) A polymerization process comprising contacting one or more 32. isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor, The process-of elaim 1, wherein the one or more initiator(s) is independently selected from the group consisting of 2-chloro-2,4,4-trimethylpentane; 2-bromo-2,4,4-trimethylpentane; 2-2-chloro-2,4,4,6,6-2-bromo-2-methylpropane; chloro-2-methylpropane: 2-bromo-2,4,4,6,6-pentamethylheptane; 1-chloro-1pentamethylheptane; methylethylbenzene; 1-chloroadamantane; 1-chloroethylbenzene; 1, 4-bis(1-chloro-1methylethyl) benzene; 5-tert-butyl-1,3-bis(1-chloro-1-methylethyl) benzene; 2acetoxy-2,4,4-trimethylpentane; 2-benzoyloxy-2,4,4-trimethylpentane; 2-acetoxy-2-2-benzoyloxy-2-methylpropane; 2-acetoxy-2,4,4,6,6methylpropane; pentamethylheptane; 2-benzoyl-2,4,4,6,6-pentamethylheptane; I-acetoxy-1methylethylbenzene; 1-accotxyadamantane; 1-benzoyloxyethylbenzene; 1,4-bis(1acetoxy-i-methylethyl) benzene; 5-tert-butyl-1,3-bis(1-acetoxy-1-methylethyl) benzene; 2-methoxy-2,4,4-trimethylpentane; 2-isopropoxy-2,4,4-trimethylpentane; 2methoxy-2-methylpropane; 2-benzyloxy-2-methylpropane; 2-methoxy-2,4,4,6,6-2-isopropoxy-2,4,4,6,6-pentamethylheptane; pentamethylheptane; I-methoxy-1methylethylbenzene; 1-methoxyadamantane; 1-methoxyethylbenzene; 1,4-bis(1-5-tert-butyl-1,3-bis(1-methoxy-1-methylethyl) methoxy-1-methylethyl) benzene; benzene, and 1,3,5-tris(1-chloro-1-methylethyl) benzene.

- 33. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of claim 1, wherein the one or more initiator(s) further comprise a weakly-coordinating anion.
- 34. (Previously presented) The process of claim 1, wherein the process is substantially absent of water.
- 35. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of claim 1, wherein the one or more initiator(s) comprise greater than 30 ppm water (based upon weight).
- 36. (Previously presented) The process of claim 1, wherein the one or more monomer(s) is independently selected from the group consisting of olefins, alpha-olefins, disubstituted olefins, isoolefins, conjugated dienes, non-conjugated dienes, styrenics, substituted styrenics, and vinyl ethers.
- 37. (Currently amended) The process of claim 1 36, wherein the one or more monomer(s) is independently selected from the group consisting of styrene, paraalkylstyrene, paramethylstyrene, alpha-methyl styrene, divinylbenzene, diisopropenylbenzene, isobutylene, 2-methyl-1-butene, 3-methyl-1-butene, 2-methyl-2-pentene, isoprene, butadiene, 2,3-dimethyl-1,3-butadiene, 8-pinene, myrcene, 6,6-dimethyl-fulvene, hexadiene, cyclopentadiene, methyl cyclopentadiene, piperylene, methyl vinyl ether, ethyl vinyl ether, and isobutyl vinyl ether.
- 38. (Previously presented) The process of claim 1, wherein the reactor is independently selected from the group consisting of a continuous flow stirred tank reactor, a plug flow reactor, a moving belt or drum reactor, a jet or nozzle reactor, a tubular reactor, a batch reactor, and an autorefrigerated boiling-pool reactor.

- 39. (Previously presented) The polymerization process of claim 1, wherein the diluent has a dielectric constant greater than 10 at -85°C.
- 40. (Original) The process of claim 39, wherein the dielectric constant is greater than 20 at -85°C.
- 41. (Original) The process of claim 40, wherein the dielectric constant is greater than 25 at -85°C.
- 42. (Currently amended) A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor. The process of claim 41, wherein the diluent has a dielectric constant is greater than 40 at -85°C.
- 43. (Previously presented) The polymerization process of claim 1, wherein polymerization process forms a polymer having a diluent mass uptake of less than 4 wt%.
- 44. (Original) The process of claim 43, wherein the polymer has a diluent mass uptake of less than 3 wt%.
- 45. (Original) The process of claim 44, wherein the polymer has a diluent mass uptake of less than 2 wt%.
- 46. (Original) The process of claim 45, wherein the polymer has a diluent mass uptake of less than 1 wt%.
- 47. (Original) The process of claim 46, wherein the polymer has a diluent mass uptake of less than 0.5 wt%.

- 48. (Previously presented) The polymerization process of claim 1, the diluent comprising methyl chloride and one or more hydrofluorocarbon(s) independently selected from the group consisting of difluoromethane, 1,1-difluoroethane, and 1,1,1,2-tetrafluoroethane.
- 49. (Original) The process of claim 48 wherein the diluent further comprises a non-reactive olefin, one or more other hydrocarbons, and/or an inert gas.
- 50. (Previously presented) The polymerization process of claim 1, the process comprising the steps of:
  reacting the one or more monomer(s) in the presence of one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s)
  (HFC's); and
  withdrawing the polymer from the reactor.
- 51. (Previously presented) The polymerization process of claim 1, the process comprising the steps of:
  - (a) introducing one or more monomer(s) into a reactor;
  - (b) adding one or more Lewis acid(s) and one or more initiator(s);
  - (c) introducing a diluent comprising one or more hydrofluorocarbon(s) (HFC's); and
  - (d) withdrawing a polymer from the reactor.
- 52. (Currently Amended) The polymerization process of claim 1 in which particles of polymer are polymerized produced using a catalyst system and a diluent comprising one or more hydrofluorocarbon(s) (HFC's).
- 53. (Currently Amended) The polymerization process of claim I in which particles of polymer are polymerized produced in a slurry using a carbocationic catalyst system and a diluent comprising one or more hydrofluorocarbon(s) (HFC's).

Page 16 of 33

- 54. (Currently Amended) The polymerization process of claim 1 in which particles of polymer are polymerized produced using a catalyst system and a diluent, the diluent having a dielectric constant greater than 18.50 at -85°C.
- 55. (Original) The process of claim 54, wherein the dielectric constant is greater than 19.00 at -85°C.
- 56. (Original) The process of claim 55, wherein the dielectric constant is greater than 20.00 at -85°C.
- 57. (Original) The process of claim 56, wherein the dielectric constant is greater than 25.00 at -85°C.
- 58. (Original) The process of claim 57, wherein the dielectric constant is greater than 30.00 at -85°C.
- 59. (Original) The process of claim 58, wherein the dielectric constant is greater than 35.00 at -85°C.
- 60. (Currently amended) The process of claim 59, wherein the dielectric constant is greater than 40.00 at -85°C. A polymerization process comprising contacting one or more isoolefins, one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor, in which particles of polymer are produced using a catalyst system and a diluent, the diluent having a dielectric constant greater than 40.00 at -85°C.
- 61. (Previously presented) The process of claim 1, wherein the diluent is independently selected from the group consisting of at least one of a hydrofluorocarbon or a blend of a hydrofluorocarbon and methyl chloride.

- 62. (Currently Amended) A polymerization process in which particles of polymer are polymerized produced using a diluent comprising one or more hydrofluorocarbon(s) (HFC's).
- ·63. (Cancelled)
- 64. (Original) A polymerization medium suitable to polymerize one or more monomer(s) to form a polymer, the polymerization medium comprising one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's).
- 65. (Cancelled)
- 66. (Cancelled)
- 67. (Cancelled)
- 68. (Cancelled)
- 69. (Cancelled)
- 70. (New) The polymerization process of claim 1, wherein the diluent has a dielectric constant greater than 40 at -85°C.
- 71. (New) A polymerization process comprising contacting one or more monomers comprising isoolefin(s), one or more Lewis acid(s), one or more initiator(s), and a diluent comprising one or more hydrofluorocarbon(s) (HFC's) in a reactor.
- 72. (New) The process of claim 71, wherein one or more hydrofluorocarbon(s) is represented by the formula:  $C_xH_yF_z$  wherein x is an integer from 1 to 40 and y and z are integers of one or more.

- 73. (New) The process of claim 72, wherein x is from 1 to 10.
- 74. (New) The process of claim 72, wherein x is from 1 to 6.
- 75. (New) The process of claim 72, wherein x is from 1 to 3.
- (New) The process of claim 71, wherein the one or more hydrofluorocarbon(s) is 76. independently selected from the group consisting of fluoromethane; difluoromethane; fluoroethane; 1,1-difluoroethane; 1,2-difluoroethane; 1,1,1trifluoromethane; 1,1,1,2-tetrafluoroethane; 1,1,2,2-1,1,2-trifluoroethane; trifluoroethane; tetrafluoroethane; 1,1,1,2,2-pentafluoroethane; 1-fluoropropane; 2-fluoropropane; 1,1difluoropropane; 1,2-difluoropropane; 1,3-difluoropropane; 2,2-difluoropropane; 1,1,3-trifluoropropane; 1,2,2-1,1,1-trifluoropropane; 1,1,2-trifluoropropane; 1,1,1,3-1,1,1,2-tetrafluoropropane; trifluoropropane; 1,2,3-trifluoropropane; tetrafluoropropane; 1,1,2,2-tetrafluoropropane; 1,1,2,3-tetrafluoropropane; 1,1,3,3-1,1,1,2,2-pentafluoropropane; tetrafluoropropane; 1,2,2,3-tetrafluoropropane; 1,1,2,2,3-1,1,1,3,3-pentafluoropropane; 1,1,1,2,3-pentafluoropropane; pentafluoropropane; 1,1,2,3,3-pentafluoropropane; 1,1,1,2,2,3-hexafluoropropane; 1,1,1,3,3,3-hexafluoropropane; 1,1,1,2,2,3,3-1,1,1,2,3,3-hexafluoropropane; 1,1,1,2,3,3,3-heptafluoropropane; 1-fluorobutane; 2heptafluoropropane; fluorobutane; 1,1-difluorobutane; 1,2-difluorobutane; 1,3-difluorobutane; 1.4difluorobutane; 2,2-difluorobutane; 2,3-difluorobutane; 1,1,1-trifluorobutane; 1,1,2trifluorobutane; 1,1,3-trifluorobutane; 1,1,4-trifluorobutane; 1,2,2-trifluorobutane; 2.2.3-trifluorobutane; 1,1,1,2-1,2,3-trifluorobutane; 1,3,3-trifluorobutane; 1,1,2,2-1,1,1,3-tetrafluorobutane; 1,1,4-tetrafluorobutane; tetrafluorobutane; 1,1,2,4-tetrafluorobutane; 1,1,3,3-1,1,2,3-tetrafluorobutane; tetrafluorobutane; 1,1,4,4-tetrafluorobutane; 1,2,2,3-1,1,3,4-tetrafluorobutane; tetrafluorobutane: 1,2,3,3-tetrafluorobutane; 1,2,3,4-1.2.2.4-tetrafluorobutane; tetrafluorobutane; tetrafluorobutane; 2,2,3,3-tetrafluorobutane; 1,1,1,2,2-pentafluorobutane; 1,1,1,2,3-1,1,1,3,3-pentafluorobutane; 1.1.1.2.4-pentafluorobutane; pentafluorobutane; 1,1,2,2,3-1,1,1,4,4-pentafluorobutane; 1,1,1,3,4-pentafluorobutane; 1,1,2,3,3-pentafluorobutane; 1,1,2,2,4-pentafluorobutane; pentafluorobutane;

> 1,1,3,3,4-pentafluorobutane; 1,2,2,3,3-1,1,2,4,4-pentafluorobutane; 1,1,1,2,2,3-hexafluorobutane; 1,2,2,3,4-pentafluorobutane; pentafluorobutane; 1,1,1,2,3,3-hexafluorobutane, 1,1,1,2,3,4-1,1,2,2,4-hexafluorobutane; 1,1,1,3,3,4-hexafluorobutane; 1,1,1,2,4,4-hexafluorobutane; hexafluorobutane: 1,1,2,2,3,3-1.1.1.4.4.4-hexafluorobutane; 1,1,1,3,4,4-hexafluorobutane; 1,1,2,2,4,4-hexafluorobutane; 1,1,2,2,3,4-hexafluorobutane; hexafluorobutane; 1,2,2,3,3,4-1,1,2,3,3,4-hexafluorobutane; 1,1,2,3,4,4-hexafluorobutane; hexafluorobutane; 1,1,1,2,2,3,3-heptafluorobutane; 1,1,1,2,2,4,4-heptafluorobutane; 1,1,1,2,3,3,4-heptafluorobutane; 1,1,1,2,3,4,4-1,1,1,2,2,3,4-heptafluorobutane; heptafluorobutane; 1,1,1,2,4,4,4-heptafluorobutane; 1,1,1,3,3,4,4-heptafluorobutane; 1,1,1,2,2,3,3,4-octafluorobutane; 1,1,1,2,2,3,4,4-octafluorobutane; 1,1,1,2,3,3,4,4octafluorobutane; 1,1,1,2,2,4,4,4-octafluorobutane; 1,1,1,2,3,4,4,4-octafluorobutane; 1,1,1,2,2,3,3,4,4-nonafluorobutane; 1,1,1,2,2,3,4,4,4-nonafluorobutane; 1-fluoro-2methylpropane; 1,1-difluoro-2-methylpropane; 1,3-difluoro-2-methylpropane; 1,1,1-1,3-difluoro-2-1,1,3-trifluoro-2-methylpropane; trifluoro-2-methylpropane; (fluoromethyl)propane; 1,1,1,3-tetrafluoro-2-methylpropane; 1,1,3,3-tetrafluoro-2-1.1.3-trifluoro-2-(fluoromethyl)propane; 1,1,1,3,3-pentafluoro-2methylpropane; methylpropane; 1,1,3,3-tetrafluoro-2-(fluoromethyl)propane; 1,1,3,4-tetrafluoro-2-1.2fluorocyclobutane; 1,1-difluorocyclobutane; (fluoromethyl)propane; 1,3-difluorocyclobutane; 1,1,2-trifluorocyclobutane; 1,1,3difluorocyclobutane; 1,1,2,2-tetrafluorocyclobutane; trifluorocyclobutane; 1,2,3-trifluorocyclobutane; 1,1,2,3,3-1,1,2,2,3-pentafluorocyclobutane; 1,1,3,3-tetrafluorocyclobutane; 1,1,2,2,3,3-hexafluorocyclobutane; 1,1,2,2,3,4pentafluorocyclobutane; 1,1,2,3,3,4-hexafluorocyclobutane; 1,1,2,2,3,3,4hexafluorocyclobutane; heptafluorocyclobutane; vinyl fluoride; 1,1-difluoroethene; 1,2-difluoroethene; 1,1,2trifluoroethene; 1-fluoropropene, 1,1-difluoropropene; 1,2-difluoropropene; 1,3difluoropropene; 2,3-difluoropropene; 3,3-difluoropropene; 1,1,2-trifluoropropene; 1,3,3-trifluoropropene; 1,1,3-trifluoropropene; 1,2,3-trifluoropropene; trifluoropropene; 3,3,3-trifluoropropene; 1-fluoro-1-butene; 2-fluoro-1-butene; 3fluoro-1-butene; 4-fluoro-1-butene; 1,1-difluoro-1-butene; 1,2-difluoro-1-butene; 1,3difluoropropene; 1,4-difluoro-1-butene; 2,3-difluoro-1-butene; 2,4-difluoro-1-butene; 3,3-difluoro-1-butene; 3,4-difluoro-1-butene; 4,4-difluoro-1-butene; 1,1,2-trifluoro-1-

> butene; 1,1,3-trifluoro-1-butene; 1,1,4-trifluoro-1-butene; 1,2,3-trifluoro-1-butene; 1,2,4-trifluoro-1-butene; 1,3,3-trifluoro-1-butene; 1,3,4-trifluoro-1-butene; 1,4,4trifluoro-1-butene; 2,3,3-trifluoro-1-butene; 2,3,4-trifluoro-1-butene; 2,4,4-trifluoro-1butene; 3,3,4-trifluoro-1-butene; 3,4,4-trifluoro-1-butene; 4,4,4-trifluoro-1-butene; 1,1,2,4-tetrafluoro-1-butene; 1.1.3.3-tetrafluoro-1-1,1,2,3-tetrafluoro-1-butene; butene; 1,1,3,4-tetrafluoro-1-butene; 1,1,4,4-tetrafluoro-1-butene; 1,2,3,3-tetrafluoro-1,2,4,4-tetrafluoro-1-butene; 1,2,3,4-tetrafluoro-1-butene; 1,4,4,4-tetrafluoro-1-butene; 1,3,4,4-tetrafluoro-1-butene; tetrasluoro-1-butene; 2,4,4,4-tetrafluoro-1-2,3,4,4-tetrafluoro-1-butene; 2.3,3,4-tetrafluoro-1-butene; 1,1,2,3,3-3.4.4.4-tetrafluoro-1-butene; 3,3,4,4-tetrafluoro-1-butene; butene; pentafluoro-1-butene; 1,1,2,3,4-pentafluoro-1-butene; 1,1,2,4,4-pentafluoro-1-butene; 1,1,3,3,4-pentafluoro-1-butene; 1,1,3,4,4-pentafluoro-1-butene; 1,1,4,4,4-pentafluoro-1-butene; 1,2,3,3,4-pentafluoro-1-butene; 1,2,3,4,4-pentafluoro-1-butene; 1,2,4,4,4pentafluoro-1-butene; 2,3,3,4,4-pentafluoro-1-butene; 2,3,4,4,4-pentafluoro-1-butene; 1,1,2,3,4,4-1,1,2,3,3,4-hexafluoro-1-butene; 3,3,4,4,4-pentafluoro-1-butene; 1,2,3,3,4,4-hexafluoro-1-1.1.2.4.4.4-hexafluoro-1-butene; hexafluoro-1-butene: 1,2,3,4,4,4-hexafluoro-1-butene; 2,3,3,4,4,4-hexafluoro-1-butene; butene; 1,1,2,3,3,4,4-heptafluoro-1-butene; 1,1,2,3,4,4,4-heptafluoro-1-butene; 1,1,3,3,4,4,4heptafluoro-1-butene; 1,2,3,3,4,4,4-heptafluoro-1-butene; 1-fluoro-2-butene; 2-fluoro-2-butene; 1,1-difluoro-2-butene; 1,2-difluoro-2-butene; 1,3-difluoro-2-butene; 1,4difluoro-2-butene; 2,3-difluro-2-butene; 1,1,1-trifluoro-2-butene; 1,1,2-trifluoro-2butene; 1,1,3-trifluoro-2-butene; 1,1,4-trifluoro-2-butene; 1,2,3-trifluoro-2-butene; 1,2,4-trifluoro-2-butene; 1,1,1,2-tetrafluoro-2-butene; 1,1,1,3-tetrafluoro-2-butene; 1,1,2,4-tetrafluoro-2-1,1,2,3-tetrafluoro-2-butene; 1,1,1,4-tetrafluoro-2-butene; butene; 1,2,3,4-tetrafluoro-2-butene; 1,1,1,2,3-pentafluoro-2-butene; 1,1,1,2,4pentafluoro-2-butene; 1,1,1,3,4-pentafluoro-2-butene; 1,1,1,4,4-pentafluoro-2-butene; 1,1,2,4,4-pentafluoro-2-butene; 1,1,1,2,3,4-1,1,2,3,4-pentafluoro-2-butene; 1,1,1,3,4,4-hexafluoro-2-1.1.1.2.4.4-hexaflyoro-2-butene; hexafluoro-2-butene; 1,1,2,3,4,4-hexafluoro-2-butene; 1,1,4,4,4-hexafluoro-2-butene; butene: 1,1,1,2,3,4,4-heptafluoro-2-butene; 1,1,1,2,4,4,4-heptafluoro-2-butene; and mixtures thereof.

- 77. (New) The process of claim 71, wherein the one or more hydrofluorocarbon(s) is independently selected from the group consisting of fluoromethane, difluoromethane, trifluoromethane, 1,1-difluoroethane, 1,1-trifluoroethane, 1,1,2-tetrafluoroethane, and mixtures thereof.
- 78. (New) The process of claim 71, wherein the diluent comprises from 15 to 100 volume % HFC based upon the total volume of the diluent.
- 79. (New) The process of claim 78, wherein the diluent comprises from 20 to 100 volume % HFC based upon the total volume of the diluent.
- 80. (New) The process of claim 79, wherein the diluent comprises from 25 to 100 volume % HFC based upon the total volume of the diluent.
- 81. (New) The process of claim 71, wherein the diluent further comprises a hydrocarbon, a non-reactive olefin, and/or an inert gas.
- 82. (New) The process of claim 71, wherein the diluent further comprises a halogenated hydrocarbon other than an HFC.
- 83. (New) The process of claim 71, wherein the diluent further comprises methyl chloride.
- 84. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>4</sub>; wherein M is a Group 4, 5, or 14 metal; and each X is a halogen.
- 85. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MR<sub>n</sub>X<sub>4-n</sub>; wherein M is Group 4, 5, or 14 metal;

each R is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4; and each X is a halogen.

86. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula M(RO)<sub>n</sub>R'<sub>m</sub>X<sub>4-(m+n);</sub> wherein M is Group 4, 5, or 14 metal; each RO is a monovalent C<sub>1</sub> to C<sub>30</sub> hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, arylalkoxy, alkylaryloxy radicals; each R' is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from

the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4;

m is an integer from 0 to 4, wherein the sum of n and m is not more than 4; and each X is a halogen.

87. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula  $M(RC=OO)_nR'_mX_{4-(m+n)}$ ;

wherein M is Group 4, 5, or 14 metal;

each RC=OO is a monovalent  $C_2$  to  $C_{30}$  hydrocarbacyl radical independently selected from the group consisting of an alkacyloxy, arylacyloxy, arylacyloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4;

m is an integer from 0 to 4, wherein the sum of n and m is not more than 4; and each X is a halogen.

88. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MOX<sub>3</sub>; wherein M is a Group 5 metal; and each X is a halogen.

- 89. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>3</sub>; wherein M is a Group 13 metal; and each X is a halogen.
- 90. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MR<sub>n</sub>X<sub>3-n</sub>: wherein M is a Group 13 metal; each R is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 1 to 3; and each X is a halogen.
- 91. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula M(RO)<sub>n</sub>R'<sub>m</sub>X<sub>3-(m+n)</sub>:
  wherein M is a Group 13 metal;
  each RO is a monovalent C<sub>1</sub> to C<sub>30</sub> hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, arylalkoxy, alkylaryloxy radicals;
  each R' is a monovalent C<sub>1</sub> to C<sub>12</sub> hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals;
  n is an integer from 0 to 3;
  m is an integer from 0 to 3, wherein the sum of n and m is from 1 to 3; and each X is a halogen.
- 92. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula M(RC=OO)<sub>n</sub>R'<sub>m</sub>X<sub>3-(m+n)</sub>; wherein M is a Group 13 metal; each RC=OO is a monovalent hydrocarbacyl radical independently selected from the group independently selected from the C<sub>2</sub> to C<sub>30</sub> group consisting of an alkacyloxy, arylacyloxy, arylacyloxy, alkylarylacyloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 3; m is a integer from 0 to 3, wherein the sum of n and m is from 1 to 3; and each X is a halogen.

- 93. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula MX<sub>y</sub>, wherein M is a Group 15 metal; each X is a halogen; and y is 3, 4 or 5.
- 94. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula:

 $MR_nX_{y-n}$ 

wherein M is a Group 15 metal;

each R is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4;

y is 3, 4 or 5, wherein n is less than y; wherein the sum of n and m is less than y each X is a halogen.

95. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula  $M(RO)_n R'_m X_{y \cdot (m+n)}$ :

wherein M is a Group 15 metal,

each RO is a monovalent  $C_1$  to  $C_{30}$  hydrocarboxy radical independently selected from the group consisting of an alkoxy, aryloxy, arylalkoxy, alkylaryloxy radicals; each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals; n is an integer from 0 to 4;

m is an integer from 0 to 4;

 $\nu$  is 3, 4 or 5, wherein the sum of n and m is less than y; and

each X is a halogen.

96. (New) The process of claim 71, wherein the one or more Lewis acid(s) is represented by the formula M(RC=OO)<sub>0</sub>R'<sub>m</sub>X<sub>y-(m+n)</sub>;

wherein M is a Group 15 metal;

each RC=OO is a monovalent  $C_2$  to  $C_{30}$  hydrocarbacyloxy radical independently selected from the group consisting of an alkacyloxy, arylacyloxy, arylacyloxy radicals;

each R' is a monovalent  $C_1$  to  $C_{12}$  hydrocarbon radical independently selected from the group consisting of an alkyl, aryl, arylalkyl, alkylaryl and cycloalkyl radicals;

n is an integer from 0 to 4;

m is an integer from 0 to 4;

y is 3, 4 or 5, wherein the sum of n and m is less than y; and each X is a halogen.

(New) The process of claim 71, wherein the one or more Lewis acid(s) is 97. independently selected from the group consisting of titanium tetrachloride, titanium tetrabromide, vanadium tetrachloride, tin tetrachloride, zirconium tetrachloride, titanium bromide trichloride, titanium dibromide dichloride, vanadium bromide trichloride, tin chloride trifluoride, benzyltitanium trichloride, dibenzyltitanium dibenzylzirconium dibromide, trichloride, benzylzirconium dichloride, methyltitanium trichloride, dimethyltitanium difluoride, dimethyltin dichloride, trichloride, n-butoxytitanium trichloride, methoxytitanium phenylvanadium tribromide, di(isopropoxy)titanium dichloride, phenoxytitanium trichloride, phenylmethoxyzirconium trifluoride, methyl methoxytitanium dichloride, methyl methoxytin dichloride, benzyl isopropoxyvanadium dichloride, acetoxytitanium benzoyloxytitanium trifluoride, tribromide. trichloride, benzoylzirconium dichloride. benzyl methyl acetoxytitanium trichloride, isopropoyloxytin benzoyloxyvanadium chloride, vanadium oxytrichloride, aluminum trichloride, boron trifluoride, gallium trichloride, indium trifluoride, ethylaluminum dichloride, methylaluminum dichloride, benzylaluminum dichloride, isobutylgallium dichloride, ethylaluminum dimethylaluminum chloride. diethylaluminum chloride,

> trimethylaluminum, sesquichloride methylaluminum sesquichloride, triethylaluminum, methoxyaluminum dichloride, ethoxyaluminum dichloride, 2,6-ditert-butylphenoxyaluminum dichloride, methoxy methylaluminum chloride, 2,6-ditert-butylphcnoxy methylaluminum chloride, isopropoxygallium dichloride, phenoxy dichloride, benzoyloxyaluminum acetoxyaluminum methylindium fluoride. dibromide, benzoyloxygallium difluoride, methyl acetoxyaluminum chloride, isopropoyloxyindium trichloride, antimony hexachloride, antimony hexafluoride, arsenic pentafluoride, antimony chloride pentafluoride, arsenic trifluoride, bismuth chlaride, tetrachloride, tetraphenylantimony trichloride arsenic fluoride tetrachloromethoxyantimony, dichloride, triphenylantimony dichloromethoxyarsine, chlorodimethoxyarsine, dimethoxytrichloroantimony, difluoromethoxyarsine, acetatotetrachloroantimony, (benzoato) tetrachloroantimony, and bismuth acetate chloride.

- 98. (New) The process of claim 71, wherein the one or more Lewis acid(s) is independently selected from the group consisting of aluminum trichloride, aluminum tribromide, ethylaluminum dichloride, ethylaluminum sesquichloride, diethylaluminum chloride, methylaluminum dichloride, methylaluminum sesquichloride, dimethylaluminum chloride, boron trifluoride, and titanium tetrachloride.
- 99. (New) The process of claim 71, wherein the one or more initiator(s) comprise a hydrogen halide, a carboxylic acid, a carboxylic acid halide, a sulfonic acid, an alcohol, a phenol, a polymeric halide, a tertiary alkyl halide, a tertiary aralkyl halide, a tertiary aralkyl ester, a tertiary aralkyl ester, a tertiary aralkyl ether, an alkyl halide, an aryl halide, an alkylaryl halide or an arylalkylacid halide.
- 100. (New) The process of claim 71, wherein the one or more initiator(s) is independently selected from the group consisting of HCl, H<sub>2</sub>O, methanol, (CH<sub>3</sub>)<sub>3</sub>CCl, C<sub>6</sub>H<sub>5</sub>C(CH<sub>3</sub>)<sub>2</sub>Cl, (2-Chloro-2,4,4-trimethylpentane) and 2-chloro-2-methylpropane.

- (New) The process of claim 71, wherein the one or more initiator(s) is independently 101. selected from the group consisting of hydrogen chloride, hydrogen bromide, hydrogen · iodide, acetic acid, propanoic acid, butanoic acid; cinnamic acid, benzoic acid, 1chloroacetic acid, dichloroacetic acid, trichloroacetic acid, trifluoroacetic acid, pchlorobenzoic acid, p-fluorobenzoic acid, acetyl chloride, acetyl bromide, cinnamyl bromide. trichloroacetylchloride, chloride, benzoyl chloride, benzoyl methanesulfonic acid. p-fluorobenzoylchloride, trifluoroacetylchloride, trifluoromethanesulfonic acid, trichloromethanesulfonic acid, p-tolucnesulfonic acid, methanesulfonyl chloride, methanesulfonyl bromide, trichloromethanesulfonyl chloride, trifluoromethanesulfonyl chloride, p-toluenesulfonyl chloride, methanol, ethanol, propanol, 2-propanol, 2-methylpropan-2-ol, cyclohexanol, benzyl alcohol, phenol, 2-methylphenol, 2,6-dimethylphenol, p-chlorophenol, p-fluorophenol, 2,3,4,5,6-pentafluorophenol, and 2-hydroxynaphthalene.
- (New) The process of claim 71, wherein the one or more initiator(s) is independently 102. selected from the group consisting of 2-chloro-2,4,4-trimethylpentane; 2-bromo-2,4,4trimethylpentane; 2-chloro-2-methylpropane; 2-bromo-2-methylpropane; 2-chloro-2,4,4,6,6-pentamethylheptane; 2-bromo-2,4,4,6,6-pentamethylheptane; 1-chloro-1methylethylbenzene; 1-chloroadamantane; 1-chloroethylbenzene; 1, 4-bis(1-chloro-1methylethyl) benzene; 5-tert-butyl-1,3-bis(1-chloro-1-methylethyl) benzene; 2acetoxy-2,4,4-trimethylpentane; 2-benzoyloxy-2,4,4-trimethylpentane; 2-acetoxy-2-2-acetoxy-2,4,4,6,6-2-benzoyloxy-2-methylpropane; methylpropane; 2-benzoyl-2,4,4,6,6-pentamethylheptane; 1-acetoxy-1pentamethylheptane; methylethylbenzene; 1-aceotxyadamantane; 1-benzoyloxyethylbenzene; 1,4-bis(1-5-tert-butyl-1,3-bis(1-acetoxy-1-methylethyl) benzene: acetoxy-I-methylethyl) benzene; 2-methoxy-2,4,4-trimethylpentane; 2-isopropoxy-2,4,4-trimethylpentane; 2methoxy-2-methylpropane; 2-henzyloxy-2-methylpropane; 2-methoxy-2,4,4,6,6-2-isopropoxy-2,4,4,6,6-pentamethylheptane; 1-methoxy-1pentamethylheptane; 1-methoxyadamantane; 1-methoxyethylbenzene; 1,4-bis(1methylethylbenzene; benzene; 5-tert-butyl-1,3-bis(1-methoxy-1-methylethyl) methoxy-1-methylethyl) benzene, and 1,3,5-tris(1-chloro-1-methylethyl) benzene.

- 103. (New) The process of claim 71, wherein the one or more initiator(s) further comprise a weakly-coordinating anion.
- 104. (New) The process of claim 71, wherein the process is substantially absent of water.
- 105. (New) The process of claim 71, wherein the one or more initiator(s) comprise greater than 30 ppm water (based upon weight).
- 106. (New) The process of claim 1, wherein the one or more monomer(s) comprises isobutylene.
- 107. (New) The process of claim 71, wherein the one or more monomer(s) comprises isobutylene and para-alkylstyrene or isoprene.
- 108. (New) The process of claim 71, wherein the reactor is independently selected from the group consisting of a continuous flow stirred tank reactor, a plug flow reactor, a moving belt or drum reactor, a jet or nozzle reactor, a tubular reactor, a batch reactor, and an autorefrigerated boiling-pool reactor.
- 109. (New) The polymerization process of claim 71, wherein the diluent has a dielectric constant greater than 10 at -85°C.
- 110. (New) The polymerization process of claim 71, wherein polymerization process forms a polymer having a diluent mass uptake of less than 4 wt%.
- 111. (New) The polymerization process of claim 71, the diluent comprising methyl chloride and one or more hydrofluorocarbon(s) independently selected from the group consisting of difluoromethane, 1,1-difluoroethane, and 1,1,2-tetrafluoroethane.
- 112. (New) The process of claim 111 wherein the diluent further comprises a non-reactive olefin, one or more other hydrocarbons, and/or an inert gas.

113. (New) The process of claim 71, wherein the diluent is independently selected from the group consisting of at least one of a hydrofluorocarbon or a blend of a hydrofluorocarbon and methyl chloride.